

# DhhP, a Cyclic di-AMP Phosphodiesterase of *Borrelia burgdorferi*, Is Essential for Cell Growth and Virulence

Meiping Ye,<sup>a,b</sup> Jun-Jie Zhang,<sup>b,c</sup> Xin Fang,<sup>d</sup> Gavin B. Lawlis,<sup>d</sup> Bryan Troxell,<sup>b</sup> Yan Zhou,<sup>a</sup> Mark Gomelsky,<sup>d</sup> Yongliang Lou,<sup>a</sup> X. Frank Yang<sup>a,b</sup>

School of Laboratory Medicine, Wenzhou Medical University, Zhejiang, China<sup>a</sup>; Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis, Indiana, USA<sup>b</sup>; Environmental Microbiology Group, Key Laboratory of Agricultural and Environmental Microbiology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, China<sup>c</sup>; Department of Biology, University of Wyoming, Laramie, Wyoming, USA<sup>d</sup>

Cyclic di-AMP (c-di-AMP) is a recently discovered second messenger in bacteria. Most of work on c-di-AMP signaling has been done in Gram-positive bacteria, firmicutes, and actinobacteria, where c-di-AMP signaling pathways affect potassium transport, cell wall structure, and antibiotic resistance. Little is known about c-di-AMP signaling in other bacteria. *Borrelia burgdorferi*, the causative agent of Lyme disease, is a spirochete that has a Gram-negative dual membrane. In this study, we demonstrated that *B. burgdorferi* BB0619, a DHH-DHHA1 domain protein (herein designated DhhP), functions as c-di-AMP phosphodiesterase. Recombinant DhhP hydrolyzed c-di-AMP to pApA in a Mn<sup>2+</sup>- or Mg<sup>2+</sup>-dependent manner. In contrast to c-di-AMP phosphodiesterases reported thus far, DhhP appears to be essential for *B. burgdorferi* growth both *in vitro* and in the mammalian host. Inactivation of the chromosomal *dhhP* gene could be achieved only in the presence of a plasmid-encoded inducible *dhhP* gene. The conditional *dhhP* mutant had a dramatic increase in intracellular c-di-AMP level in comparison to the isogenic wild-type strain. Unlike what has been observed in Gram-positive bacteria, elevated cellular c-di-AMP in *B. burgdorferi* did not result in an increased resistance to  $\beta$ -lactamase antibiotics, suggesting that c-di-AMP's functions in spirochetes differ from those in Gram-positive bacteria. In addition, the *dhhP* mutant was defective in induction of the  $\sigma^S$  factor, RpoS, and the RpoS-dependent outer membrane virulence factor OspC, which uncovers an important role of c-di-AMP in *B. burgdorferi* virulence.

Nucleotide signaling molecules play a key role in regulating cellular activities throughout all kingdoms of life. Cyclic di-AMP (c-di-AMP) is a second messenger recently identified in *Bacteria* (for reviews, see references 1, 2, 3, 4, and 5). It was initially discovered in a structural study of the DisA protein (DNA integrity scanning protein) of *Bacillus subtilis* and *Thermotoga maritima*, where it was found bound to DisA (6). Cyclic di-AMP is synthesized from two molecules of ATP by diadenylate cyclase (DAC), which contains a conserved DisA\_N domain (diadenylate cyclase, DUF147), and is hydrolyzed to pApA, or AMP, by phosphodiesterases (PDEs) containing the DHH (conserved Asp-His-His motif) and DHHA (DHH-associated domain) protein domains (5, 7). Genomic analyses revealed that proteins containing DisA\_N and DHH-DHHA domains are encoded in the genomes of many *Bacteria* as well as some *Archaea*, suggesting that c-di-AMP signaling systems are widespread (1, 5).

Although the function of c-di-AMP remains poorly understood, it appears to be an essential molecule, since attempts to inactivate the DAC genes have failed in all bacterial species where they have been made (8–14). *B. subtilis* contains three DAC genes, but at least one of the three is needed for cell growth (15). On the other hand, the c-di-AMP-specific PDE genes proved to be dispensable, and the corresponding mutants have been obtained in several Gram-positive bacteria (*Firmicutes*), including *B. subtilis*, *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Listeria monocytogenes*, and *Lactococcus lactis* (7, 8, 16–18). Most of these PDEs belong to the GdpP protein family (GGDEF domain protein-containing Phosphodiesterase) (Fig. 1), as they contain N-terminal transmembrane helices linked to the highly degenerate GGDEF domain and PAS domain (Per-Arnt-Sim domain, known to be involved in binding to small-molecule metabolites) (19), followed by the DHH-DHHA1 domain module (20).

Studies on the c-di-AMP PDE-defective mutants indicate that increased intracellular c-di-AMP levels correlate with increased cell resistance to environmental stresses, such as acid stress, heat stress, UV irradiation, and antibiotics (8, 16, 17, 21, 22). Increased stress resistance may be due to an increased peptidoglycan cross-linking (8, 15). The *gdpP* mutant of *S. aureus* also rescued the severe growth defects of mutants lacking lipoteichoic acids (21). In addition, elevated intracellular levels of c-di-AMP also resulted in impaired cell size or cell division in *B. subtilis* (5), *S. aureus*, and *S. pneumoniae* (7, 15, 21). Deletion of the c-di-AMP-dependent transcriptional repressor DarR in *Mycobacterium smegmatis* also led to increased cell length (23). Recently identified RCK\_C domain (regulator of conductance of K<sup>+</sup>) was regarded as a c-di-AMP binding domain and a c-di-AMP binding protein (CabP) in *S. pneumoniae* interacting with a K<sup>+</sup> transporter has linked c-di-AMP with K<sup>+</sup> transport (24, 25).

In term of pathogenesis, c-di-AMP has been shown to function as PAMP (pathogen-associated molecular patterns), secreted through the multidrug resistance transporter (MTR) system of *L. monocytogenes* to the host cytosol, where it triggers STING (stimulator of interferon genes)-mediated activation of the interferon

Received 9 January 2014 Returned for modification 29 January 2014

Accepted 11 February 2014

Published ahead of print 24 February 2014

Editor: A. J. Bäuml

Address correspondence to Yongliang Lou, ly093@gmail.com, or X. Frank Yang, xfyang@iupui.edu.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/IAI.00030-14




FIG 1 Diagram of the domain structure of the GdpP-family c-di-AMP PDEs and *B. burgdorferi* DhhP. TM, transmembrane domain.

(IFN) response. Inactivation of c-di-AMP PDE genes resulted in reduced infection in several pathogens, including *L. monocytogenes*, *S. pyogenes*, and *S. pneumoniae* (7, 8, 22, 26).

Virtually all the studies on c-di-AMP done thus far were conducted in Gram-positive bacteria, with the exception of one recent study showing that the Gram-negative bacterium *Chlamydia trachomatis* produces c-di-AMP (27). It was shown that c-di-AMP produced by *C. trachomatis* DAC is largely responsible for *Chlamydia*-induced STING-dependent type I IFN response in infected mammalian cells. However, the cellular functions of c-di-AMP in Gram-negative bacteria have not yet been examined.

*Borrelia burgdorferi* is a spirochetal pathogen and the causative agent of Lyme disease (28). Spirochetes are a phylogenetically distinct branch of *Bacteria*. Similar to other Gram-negative bacteria, spirochetes have a dual membrane structure with a thin layer of peptidoglycan located in the periplasmic space. The natural hosts of *B. burgdorferi* are ticks and mammals. *B. burgdorferi* undergoes massive transcriptome and proteome changes upon pathogen migration between these two drastically different hosts (29–31). Amazingly, *B. burgdorferi* accomplishes these dramatic processes with its streamlined genome and limited regulatory repertoire (32). For example, in contrast to many bacteria that have numerous c-di-GMP signaling systems (33–36), *B. burgdorferi* has only one c-di-GMP synthase (BB0419, diguanylate cyclase), and it plays an essential role in spirochetal survival in ticks as well as modulating *B. burgdorferi* motility and mammalian infection (37–43).

In this study, we began investigating the c-di-AMP signaling pathways of *B. burgdorferi*. We identified a gene for the DHH-DHHA1 domain protein BB0619, herein designated DhhP, in the *B. burgdorferi* genome and demonstrated that the protein functions as a c-di-AMP phosphodiesterase. Surprisingly, we found that, unlike what has been reported for Gram-positive bacteria, DhhP is essential for *B. burgdorferi* growth, and conditional inactivation of *dhhP* did not result in an increased resistance to  $\beta$ -lactam antibiotics, suggesting that c-di-AMP functions may differ in different phyla of bacteria. In addition, the conditional *dhhP* mutant is also defective in production of the major virulence factor OspC and in mammalian infection.

## MATERIALS AND METHODS

**Bacterial strains and culture conditions.** The low-passage, virulent *B. burgdorferi* strain 5A4NP1 (Table 1) (a gift from H. Kawabata and S.

Norris, University of Texas Health Science Center at Houston) was derived from wild-type strain B31 by inserting a kanamycin resistance marker in the restriction modification gene *bbe02* on plasmid lp25 (44). *Borrelia* organisms were cultivated in Barbour-Stoenner-Kelly (BSK-II) medium supplemented with 6% normal rabbit serum (Pel Freez Biologicals, Rogers, AR) (45) at 37°C with 5% CO<sub>2</sub>. Relevant antibiotics were added to the cultures with the following final concentrations: 250  $\mu$ g/ml for kanamycin, 50  $\mu$ g/ml for streptomycin, and 50  $\mu$ g/ml for gentamicin. The constructed suicide vector (pMP001) was maintained in *Escherichia coli* strain DH5 $\alpha$ . The antibiotics concentration used in *E. coli* were as follows: ampicillin, 100  $\mu$ g/ml; chloramphenicol, 20  $\mu$ g/ml; and gentamicin, 10  $\mu$ g/ml.

**Protein expression and purification.** To obtain the recombinant protein DhhP-6 $\times$ His, the full-length *dhhP* (bb0619) gene was PCR amplified from *B. burgdorferi* strain B31 genomic DNA using the primer pair pLATE31-dhhP F/pLATE31-dhhP R (Table 2) and cloned into pLATE31 (Thermo Scientific aLICator LIC cloning and expression system) to generate pMP032. The DhhP-6 $\times$ His fusion protein was expressed in BL21(DE3) and purified as previously described (46). Briefly, exponentially grown cells ( $A_{600}$ , 0.6) were induced for 5 h with 0.3 mM (final concentration) isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) at 25°C. Cells were pelleted and resuspended in a buffer containing 300 mM NaCl, 50 mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.4), 5% glycerol, and protease inhibitors (phenylmethylsulfonyl fluoride and P8465; Sigma, St. Louis, MO) at the concentrations specified by the manufacturer. Cells were subjected to disruption in a French pressure minicell (Spectronic Instruments), followed by brief sonication (Sonifier 250; Branson). Crude cell extracts were centrifuged at 35,000  $\times$  g for 25 min, and the supernatant was loaded onto Co<sup>2+</sup> resin (Pierce) for affinity purification. After washing, proteins were eluted with 280 mM imidazole and desalted via Zeba Spin desalting columns (7-kDa cutoff; Pierce) that had been pre-equilibrated with buffer (300 mM NaCl, 0.5 mM EDTA, 10% glycerol, 50 mM NaH<sub>2</sub>PO<sub>4</sub> [pH 7.4]). Protein purity was assessed by SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis). Protein concentration was measured using a Bradford protein assay kit (Bio-Rad).

**Enzymatic assay.** The phosphodiesterase assay conditions were essentially as described previously (17), with minor modifications. A reaction mixture (total volume, 0.6 ml) contained 0.5  $\mu$ M DhhP and 50  $\mu$ M c-di-AMP (Axxora) in the buffer 20 mM Tris (pH 8.0), 80 mM KCl, 10 mM metal cation (2.5 mM for Mn<sup>2+</sup>). The reaction was carried out at 37°C. Aliquots were withdrawn and immediately boiled for 5 min, followed by centrifugation at 15,000  $\times$  g for 10 min. The supernatant was filtered through a 0.22- $\mu$ m filter and analyzed by reversed-phase high-pressure liquid chromatography (HPLC) (Summit HPLC system; Dionex, Sunnyvale, CA) on a 15-cm by 4.6-mm Supelcosil LC-18-T column (Sigma) (47).

**Measurements of intracellular c-di-AMP.** *B. burgdorferi* organisms were grown in BSK-II medium to stationary phase, harvested in 4°C at 12,000  $\times$  g for 10 min, and immediately frozen in liquid nitrogen before being stored at  $-80^{\circ}\text{C}$ . Nucleotides were extracted from frozen *B. burgdorferi* biomass by a mixture of 40% methanol–40% acetonitrile in 0.1 N formic acid as described earlier (48). The nucleotides were separated by HPLC and quantified by using liquid chromatography-tandem mass spectrometry (LC-MS/MS) on a Quattro Premier XE mass spectrometer (Waters Corporation) coupled with an Acquity Ultra Performance LC system (Waters Corporation) at the Michigan State Mass Spectrometry Core Facility. Cyclic di-AMP standards of known concentrations were

TABLE 1 *B. burgdorferi* strains used in this study

Strain	Description	Reference or source
5A4NP1	<i>B. burgdorferi</i> B31 with <i>bbe02</i> disrupted with a kanamycin-resistant marker	44
5A4NP1/pJJ01	5A4NP1 harboring an extra copy of the IPTG-inducible gene <i>dhhP</i>	This study
5A4NP1 <i>dhhP</i> /pJJ01	5A4NP1/pJJ01 with the endogenous gene <i>dhhP</i> disrupted by a gentamicin-resistant marker	This study

TABLE 2 Primers used in the study

Primer	Sequence (5' → 3')	Purpose
pLATE31-dhhP F	AGAAGGAGATATAACTATGAGAGATGTTATTAATTTTATT	Purify DhhP
pLATE31-dhhP R	GTGGTGGTGATGGTGATGGCCTAAATAAATATTATCCTTAATG	Purify DhhP
bb0619-UF	TTGCGAATTCTCCCAAGCTTAAAAGCTCTTCTTGT	Construct pMP001
bb0619-UR	TTCGGACGTCTCAAAATCAGGATCTTTGTGCCC	Construct pMP001
bb0619-DF	TCGAGACGTCAGATTTTGGAGGCGGTGGGCA	Construct pMP001
bb0619-DR	TAGCGGATCCAAGGGGGATTGGCGCAGCAC	Construct pMP001
pJJ01-F	CCATGGAGCTCTTTTGCAGGAAGGTTTGCAGAA	Construct pJJ01
pJJ01-R	GGCCTCGAGTAAATAAATATTATCCTTAATGTATG	Construct pJJ01
dhhP ID-P1	ATGTTACGCAGCAGCAACGATGTTACGC	Verify the dhhP mutant
dhhP ID-P2	CTCCAGGATCCGTCGACTTTAGTGGCGGTACTTGGGTGCG	Verify the dhhP mutant
dhhP ID-P3	ATTTCTGTGCTTTTCTGCCAT	Verify the dhhP mutant
dhhP ID-P4	TGGAACAACACCTTGGCAT	Verify the dhhP mutant
dhhP ID-P5	AGTCGGCCATTACGTGGAAG	Verify the dhhP mutant
dhhP ID-P6	ATCAAGAAGGGGGATTGGCG	Verify the dhhP mutant
dhhP ID-P7	TTCGGAGACGTAGCCACCTA	Verify the dhhP mutant
dhhP ID-P8	CAACAACCGTCTTGGTTCG	Verify the dhhP mutant
dhhP ID-P9	AGAAGGAGATATAACTATGAGAGATGTTATTAATTTTATT	Verify the dhhP mutant
dhhP ID-P10	GTGGTGGTGATGGTGATGGCCTAAATAAATATTATCCTTAATG	Verify the dhhP mutant
qdhP-F	GGGCTATATTGATCCTTTTGCACC	qRT-PCR
qdhP-R	TGAAACCAGTCTTGCAACCA	qRT-PCR
qbosR-F	ACTGTTGCTTTTGATAGGCTTGG	qRT-PCR
qbosR-R	CGCATTGGAAGAAAGTCGGCA	qRT-PCR

used for quantification. For intracellular c-di-AMP concentration measurements, the following simplifying assumptions were made: a *Borrelia* cell has a cylindrical shape 0.25  $\mu\text{m}$  in diameter and 25  $\mu\text{m}$  in length (49); therefore, an average cell volume was calculated to be  $1.227 \times 10^{-15}$  liter.

**Construction of the conditional *dhhP* mutant.** We first constructed a wild-type strain carrying a plasmid-borne copy of *dhhP* under the control of an IPTG-inducible promoter. First, the full-length *dhhP* gene was PCR amplified from 5A4NP1 genomic DNA using the primer pair pJJ01-F/pJJ01-R and cloned into an IPTG-inducible shuttle vector, pJJ275, a derivative of pJSB275 (50) with the addition of a hemagglutinin (HA) epitope tag (amino acid sequence, YPYDVPDYA) (51) at the C terminus of the inserted gene. The resulting plasmid was designed pJJ01. pJJ01 was then transformed into 5A4NP1 using the method described previously (52, 53). The final strain, 5A4NP1/pJJ01, was confirmed by immunoblotting with anti-HA antibody.

To inactivate the endogenous *dhhP* gene in 5A4NP1/pJJ01, a suicide vector, pMP001, was constructed for homologous recombination as following: the regions of DNA corresponding to 1.4 kb upstream and 1.4 kb downstream of *dhhP* were PCR amplified from 5A4NP1 genomic DNA with the primer pairs bb0619-UF/bb0619-UR and bb0619-DF/bb0619-DR (Table 2), respectively. The resulting PCR fragments were cloned into the vector pSC-A-amp/kan (Agilent Technologies) by T-A cloning and confirmed by restriction enzyme digestion and sequencing. The *aacI* gentamicin-resistant marker driven by the *B. burgdorferi* *flaB* promoter (54) was digested out from pMH85R at the *AatII* sites (38) and was ligated between the upstream and downstream fragments of *dhhP* to generate the suicide vector pMP001. pMP001 was confirmed by both restriction enzyme digestion and sequencing and used for transformation. Cell lysates of positive *Borrelia* transformants were analyzed by PCR to confirm corrected marker insertion and inactivation of the chromosomal copy of *dhhP*. Plasmid profiles of the confirmed 5A4NP1 *dhhP*/pJJ01 clones were determined by multiple PCR analyses with 19 pairs of primers specific for each of the endogenous plasmids as reported by Bunikis et al. (55). One of the 5A4NP1 *dhhP*/pJJ01 clones that had plasmid profiles identical to those of the parental strain (5A4NP1) was chosen for further study.


**Growth curve and cell morphology.** *B. burgdorferi* strains, including the conditional *dhhP* mutant, were first grown in the presence of 1 mM

IPTG. To avoid carryover of IPTG from the seed cultures used for inoculation, harvested cells were washed twice with PBS to remove residual IPTG. Then the IPTG-free cells were recultured in the fresh BSK-II medium with (5A4NP1 *dhhP*/pJJ01+IPTG) or without (5A4NP1 *dhhP*/pJJ01–IPTG) IPTG. Either a low ( $3 \times 10^4$  cells/ml) or high ( $3 \times 10^5$  cells/ml) dose of cells was used for inoculation (indicated in the figure legends). Cell numbers were enumerated in every 12 or 24 h under dark-field microscopy. Cell morphology was visualized by dark-field microscopy, and the images were taken at a magnification of  $\times 500$ . Ten different fields were selected for measuring the cell length of each strain, and 10 spirochetes in each field were randomly selected for measurement.

**In vitro antimicrobial susceptibility test.** MICs of *B. burgdorferi* strains were determined by a standard broth microdilution method using 96-well tissue culture plates as previously described (56). Wild-type 5A4NP1, 5A4NP1 *dhhP*/pJJ01+IPTG, and 5A4NP1 *dhhP*/pJJ01–IPTG were incubated at  $10^6$  cells per ml in BSK-II medium containing a series of 2-fold dilutions of ampicillin, aztreonam, cefuroxime, cefixime, vancomycin, and oxacillin (a total volume of 220  $\mu\text{l}$  per well). The culture plates were sealed with Parafilm and were incubated for 72 h at 37°C. The lowest concentration of antibiotics that showed inhibition (lack of a color change of the BSK II medium) was interpreted as the MIC.

**Quantitative RT-PCR (qRT-PCR) and qPCR.** RNA samples were extracted from *B. burgdorferi* cultures using the RNeasy minikit (Qiagen, Valencia, CA) according to the manufacturer's protocols. Three independent culture samples were used for each strain. Digestion of contaminating genomic DNA in RNA samples was performed using RNase-free DNase I (Promega), and removal of DNA was confirmed by PCR amplification for the *B. burgdorferi* *flaB* gene. cDNA was synthesized using the SuperScript III reverse transcriptase with random primers (Invitrogen, Carlsbad, CA). To quantify the transcript levels of genes of interest, an absolute quantitation method was used by creating a standard curve in qPCR assay by following the manufacturer's protocol (Stratagene, La Jolla, CA). Briefly, the PCR product of *recA* gene served as a standard template. A series of 10-fold dilutions ( $10^2$  to  $10^8$  copies/ml) of the standard template was prepared, and qPCR was performed to generate a standard curve by plotting the initial template quantity against the cycle threshold ( $C_T$ ) values for the standards. The quantity of the targeted genes in cDNA samples was calculated using their  $C_T$  values and the standard





**FIG 2** Phosphodiesterase activity of *B. burgdorferi* DhhP *in vitro*. (A) DhhP-6×His protein following affinity purification. (B) Cyclic di-AMP hydrolysis by *B. burgdorferi* DhhP monitored by HPLC. The reaction mixture contained 0.5 μM DhhP and 50 μM c-di-AMP in the buffer (20 mM Tris, 80 mM KCl, 2.5 mM Mn<sup>2+</sup> [pH 8.0]) and was incubated at 37°C. Nucleotides were separated and analyzed using reversed-phase HPLC. Shown are nucleotide profiles before (0 min) and after (1 min incubation) addition of DhhP. (C) Metal cation dependence of c-di-AMP hydrolysis activity of DhhP. (D) pH dependence of c-di-AMP hydrolysis activity of DhhP.

curve. Samples were performed in triplicate using the ABI 7000 sequence detection system and green PCR master mix (ABI, Pleasanton, CA). Levels of the target gene transcript were reported per 100 copies of *recA* transcripts.

**Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting.** Spirochetes were harvested by centrifugation at 8,000 × g for 10 min and washed three times with PBS (pH 7.4) at 4°C. Pellets were suspended in SDS buffer containing 50 mM Tris-HCl (pH 8.0), 0.3% sodium dodecyl sulfate (SDS), and 10 mM dithiothreitol (DTT). Cell lysates (5 × 10<sup>7</sup> cells per lane) were separated by 12% SDS-polyacrylamide gel electrophoresis (PAGE) and transferred to nitrocellulose membranes (GE-Healthcare, Milwaukee, WI). Membranes were blotted with monoclonal antibodies against FlaB, OspC, RpoS, and BosR (57–59) with 1:500, 1:1,000, 1:50, and 1:500 dilutions, respectively, and then with goat anti-mouse IgG horseradish peroxidase (HRP)-conjugated secondary antibody (1:1,000; Santa Cruz Biotechnology). Detection of horseradish peroxidase activity was determined by the enhanced chemiluminescence method using luminol and peroxide as the substrate (Thermo Pierce ECL Western blotting substrate) and subsequently by exposure to X-ray film.

**Dialysis membrane chamber (DMC) implantation and mouse infection.** For cultivation of *B. burgdorferi* in the DMC mammalian host model, 5 ml of spirochetes (10<sup>3</sup> organisms per ml) in BSK-II medium were inoculated into dialysis tubing (Fisherbrand regenerated cellulose) and implanted into the peritoneal cavities of female Sprague-Dawley rats (two tubes per rat) (60, 61). For the rat implanted with 5A4NP1 *dhhP*/pJJ01 + IPTG, 10 mM IPTG was added to the drinking water (62). Dialysis tubing was harvested 14 days after implantation, and the growth of spirochetes were determined by examination of the cultures under the dark-field microscope.

For mouse infection, 3- or 4-week-old C3H/HeN mice (Harlan, Indianapolis, IN) were subcutaneously inoculated with 1 × 10<sup>5</sup> spirochetes. For mice infected with 5A4NP1 *dhhP*/pJJ01 + IPTG, 10 mM IPTG was added to the drinking water and given via intraperitoneal (IP) injection (100 μl of 10 mM IPTG) every 2 days (62). Ear punch biopsy specimens were collected 14 days after infection and cultivated in 2 ml of BSK-II medium (Sigma-Aldrich, St. Louis, MO) containing a mixture of phosphomycin (2 mg/ml), rifampin (5 mg/ml), and amphotericin B (250 mg/ml) (Sigma-Aldrich). All cultures were maintained at 37°C and examined for the presence of spirochetes by dark-field microscopy beginning 5 days after inoculation. A single growth-positive culture was used as the criterion to determine positive mouse infection.


## RESULTS

***B. burgdorferi* DhhP (BB0619) is a c-di-AMP phosphodiesterase.** The c-di-AMP-specific phosphodiesterases (PDE) studied

thus far are from Gram-positive bacteria and belong to the GdpP family (5, 7, 17, 26). GdpP often contains a DHH-DHHA1 domain together with several regulatory domains (Fig. 1). The C-terminal DHH-DHHA1 module is responsible for c-di-AMP cleavage (17). A BLAST search revealed that the *B. burgdorferi* B31 genome encodes a single DHH-DHHA1-containing protein, BB0619, that lacks other regulatory domains (Fig. 1). We designated it DhhP (DHH-DHHA1 domain protein).

To investigate whether DhhP has c-di-AMP phosphodiesterase activity, a C-terminally 6×His-tagged recombinant DhhP was expressed and purified (Fig. 2A). DhhP was then incubated with c-di-AMP in reaction buffers containing various metal cations. The result showed that DhhP hydrolyzed c-di-AMP to 5'-pApA, with the maximum activity being observed with Mn<sup>2+</sup> as a cofactor, and the optimal pH of 8.5 (Fig. 2B to D), similar to the observations made for other c-di-AMP PDEs (5, 17). Mg<sup>2+</sup> could also function as an effective metal cofactor, but Zn<sup>2+</sup> and Ca<sup>2+</sup> were inhibitory. Thus, we conclude that *B. burgdorferi* DhhP has a c-di-AMP phosphodiesterase activity.

**DhhP is essential for cell growth *in vitro*.** To investigate the function of DhhP in *B. burgdorferi*, we initially attempted to inactivate *dhhP* in wild-type *B. burgdorferi*. However, multiple attempts were unsuccessful. We reasoned that *dhhP* may be essential for *B. burgdorferi*, despite the fact that other c-di-AMP PDE genes have been readily inactivated in other bacteria (8, 16, 17, 21, 22). To test this hypothesis, we performed a conditional knockout of *dhhP* by first transforming the shuttle vector (pJJ01) carrying an IPTG-inducible *dhhP* gene into wild-type *B. burgdorferi* strain 5A4NP1, resulting in strain 5A4NP1/pJJ01 (Fig. 3A). Then the suicide vector used for inactivation of *dhhP* was transformed into 5A4NP1/pJJ01 in the presence or absence of IPTG. Ninety-three positive colonies were obtained with one independent transformation when IPTG was included in the medium, while no positive colonies were obtained with 5A4NP1 or 5A4NP1/pJJ01 – IPTG. PCR analyses confirmed that these positive clones were conditional *dhhP* mutants that contain a disrupted endogenous *dhhP* gene and an inducible *dhhP* on the shuttle vector (Fig. 3B). The endogenous plasmid profiles of the conditional *dhhP* mutant were determined using the method developed by Bunikis et al. (55). One of the *dhhP* mutant clones that had plasmid profiles identical to those of the parental strain, 5A4NP1, was chosen for further study (Fig. 3C).



**FIG 3** Construction of the conditional *dhhP* mutant. (A) Strategy for constructing the *dhhP* mutant. Arrows indicate the positions of primers (labeled above the arrows) for PCR analyses. (B) PCR analysis of the wild-type (W) and mutant (M) strains. The specific primer pairs used in PCR are indicated at the top. The arrows indicate the disrupted endogenous *dhhP* located in the chromosome (upper arrow) and an inducible copy of *dhhP* located on the shuttle vector (lower arrow). (C) Endogenous plasmid profiles of the wild-type (W) and mutant (M) strains obtained by PCR analysis using the method published by Bunikis et al. (55). cp, circular plasmid; lp, linear plasmid. Letters on each side indicate the bands corresponding to each endogenous plasmid defined previously for *B. burgdorferi* strain B31 (32).


To confirm that *dhhP* is essential for *B. burgdorferi* growth, wild-type and the conditional *dhhP* mutant strains grown in the presence of IPTG were first harvested and washed to remove IPTG. The IPTG-free cells were then inoculated into fresh BSK-II medium supplemented with various concentrations of IPTG. The result showed that the growth of the conditional *dhhP* mutant was IPTG dependent (Fig. 4A). When the initial inoculated culture concentration was high ( $3 \times 10^5$  spirochetes per ml) and the IPTG concentration was less than 10  $\mu$ M, the *dhhP* mutant replicated 5 or 6 times and then stopped (reaching about  $1 \times 10^7$  spirochetes per ml) (Fig. 4A). Supplementing the medium with  $>10 \mu$ M IPTG restored the growth of the *dhhP* mutant to the level of wild-type spirochetes. To investigate if the *dhhP* mutant could reach a density of  $10^7$  cells/ml, we inoculated the culture at a lower cell density ( $3 \times 10^4$  cells/ml). As shown in Fig. 4B, when IPTG was absent, the *dhhP* mutant replicated a few times and stopped, and it was not able to grow to  $10^7$  cells/ml.

We reasoned that the initial small number of replications observed for the conditional *dhhP* mutant observed in Fig. 4A was likely due to the presence of DhhP or IPTG in the initial cells used for inoculation. After cells had replicated several times, DhhP was depleted or degraded, and the cells then stopped replicating. To test this possibility, we used stationary-phase cells (Fig. 4A, arrow) for inoculation into fresh IPTG-free BSK-II medium at a higher


dose ( $3 \times 10^5$  per ml). As shown in Fig. 4C, the *dhhP* mutant was no longer able to replicate even for the first few times. Note that these cells remained viable under dark-field microscope (data not shown), and they replicated normally in the medium supplemented with 1 mM IPTG (Fig. 4C). Taking these data together, we conclude that DhhP is indispensable for the growth of *B. burgdorferi*.

**Deletion of *dhhP* results in an elevated cellular c-di-AMP level.** We first confirmed that the conditional *dhhP* mutant no longer expressed *dhhP* mRNA. mRNA was extracted from wild-type cells or 5A4NP1 *dhhP*/pJJ01 cells grown with and without IPTG at the stationary phase (Fig. 4A, arrow). As shown in Fig. 5A, in the absence of IPTG, the conditional mutant had no detectable level of *dhhP* transcript by RT-PCR analysis. We then subjected the cells harvested under the same conditions to measurement of the cellular concentrations of c-di-AMP. Given that DhhP was predicted to be the only DHH-DHHA-encoding protein in the *B. burgdorferi* genome, inactivation of *dhhP* was expected to result in an increased level of c-di-AMP. Indeed, the level of c-di-AMP increased more than 40-fold in the conditional *dhhP* mutant without IPTG over that in the wild-type *B. burgdorferi* strain (Fig. 5B). This result provided *in vivo* evidence that DhhP acts as a c-di-AMP PDE in *B. burgdorferi*.

**DhhP is essential for *B. burgdorferi* growth *in vivo*.** The data



**FIG 4** The *dhhP* mutant has impaired growth. (A) Wild-type strain and the conditional mutant 5A4NP1 *dhhP*/pJJ01 were cultured in BSK-II medium with 1 mM IPTG, harvested at mid-log phase, and then washed twice with fresh BSK-II medium to remove residual IPTG. The washed cells were then recultured in BSK-II medium with various concentrations of IPTG. The initial cell density was  $3 \times 10^5$  cells/ml. Spirochetes were enumerated under a dark-field microscope. Each data point is the average of data from three independent cultures. The arrow indicates the time point taken for the next experiment (C), when cells were at stationary phase. (B) Same as panel A except that the cultures were inoculated at a lower cell density ( $3 \times 10^4$  cells/ml). (C) Reinoculation of the stationary-phase cells from the culture used for panel A into fresh BSK-II medium.




**FIG 5** Inactivation of *dhhP* results in an increased intracellular c-di-AMP level. (A) Confirmation of inactivation of *dhhP* by RT-PCR analyses. Wild-type spirochetes or the conditional *dhhP* mutant grown in the absence or presence (1 mM) of IPTG were taken from cultures at day 4 (Fig. 4A). (B) Intracellular c-di-AMP concentrations. The IPTG concentration was 1 mM. Nucleotides were extracted from frozen *B. burgdorferi* organisms, separated by HPLC, and quantified by using tandem mass spectrometry as described in Materials and Methods.

above showed that the *dhhP* mutant has an impaired growth phenotype under the *in vitro* cultivation conditions. To determine whether DhhP is required for *B. burgdorferi* replication in the mammalian host environment, the wild type and the 5A4NP1 *dhhP/pJJ01* strains with and without IPTG were grown in dialysis membrane chambers (DMCs) implanted in rat peritoneal cavities, which were developed by Akins et al. (60). For the IPTG group, 10 mM IPTG was also provided in rat drinking water (62). Fourteen days after cultivation, DMCs were harvested and the cultures were examined under a dark-field microscope. As shown in Table 3, the wild type and the conditional *dhhP* mutant spirochetes supplemented with IPTG were readily detectable in DMC cultures, whereas the *dhhP* mutant without IPTG was not detected in DMCs.

We further performed mouse infection experiments for the conditional *dhhP* mutants. The wild type and the 5A4NP1 *dhhP/pJJ01* strains were needle inoculated into groups of mice ( $1 \times 10^5$  spirochetes per mouse). One group of the mice inoculated with 5A4NP1 *dhhP/pJJ01* was provided with 100  $\mu$ l 10 mM IPTG by IP every 2 days. Fourteen days after inoculation, ear punch biopsy specimens were collected from mice for cultivation of spirochetal growth. All mice inoculated with either wild-type *B. burgdorferi* or 5A4NP1 *dhhP/pJJ01* supplied with IPTG were infected. No mice

**TABLE 3** *dhhP* is required for *Borrelia* growth *in vivo*

Strain	Growth in DMC	No. of infected mice/total
5A4NP1	+	5/5
5A4NP1 <i>dhhP</i> +IPTG	+	4/4
5A4NP1 <i>dhhP</i> -IPTG	—	0/6



**FIG 6** Growth phase-dependent *dhhP* expression. Wild-type strain 5A4NP1 was grown in standard BSK-II medium at 37°C and harvested at different stages of growth. M, mid-logarithmic phase; L, late logarithmic phase; S, stationary phase. Bars represent *dhhP* mRNA levels, and black dots represent the culture doubling times. RNA was extracted and subjected to RT-PCR analysis. The data are from one representative experiment with three independent cultures. Each data point is the average from three independent cultures.

inoculated with 5A4NP1 *dhhP/pJJ01* without IPTG were infected. Taken together, these results indicate that DhhP is essential for *B. burgdorferi* replication both *in vitro* and *in vivo*.

**Growth phase-dependent expression of *dhhP*.** Given that DhhP plays an essential role in *B. burgdorferi* growth, we investigated whether *dhhP* expression correlates with the growth rate of *B. burgdorferi*. Wild-type *B. burgdorferi* 5A4NP1 spirochetes grown in standard BSK-II medium at 37°C were harvested at different growth phases and subjected to qRT-PCR analyses. As shown in Fig. 6, the *dhhP* transcript level correlated with the growth rate. Growth rate-dependent expression of *dhhP* is consistent with its essential role in *B. burgdorferi* growth.

**The *dhhP* mutant has elongated cells.** Previous reports indicated that alterations in cellular c-di-AMP were associated with changes in cell morphology. We observed that the conditional *dhhP* mutant spirochetes grown in the absence of IPTG had exhibited five times longer cells than the wild type or the 5A4NP1 *dhhP/pJJ01* spirochetes grown in the presence of IPTG (Fig. 7), suggesting that DhhP plays an important role in cell division.

**The *dhhP* mutant does not have increased resistance to  $\beta$ -lactamase antibiotics.** Increased intracellular c-di-AMP concentration has been shown to affect cell wall/peptidoglycan structure and lead to increased resistance to  $\beta$ -lactam antibiotics in several Gram-positive bacteria, including *B. subtilis*, *S. aureus*, *S. pyogenes*, and *L. monocytogenes*. To examine whether deletion of *dhhP* affects *B. burgdorferi* susceptibility to cell wall-targeting antimicrobials, MICs of the conditional *dhhP* mutant were measured using the broth microdilution method in 96-well tissue culture plates (56). The wild type and 5A4NP1 *dhhP/pJJ01* with or without IPTG were incubated with a series of dilutions of ampicillin, aztreonam, cefuroxime, cefixime, vancomycin, and oxacillin at 37°C. Note that we inoculated cells at a high dose ( $10^6$  cells per ml) so that the growth of the conditional *dhhP* mutant could be observed. MICs were calculated 72 h after incubation.

Unlike findings made with Gram-positive bacteria, in which c-di-AMP *pde* inactivation resulted in an enhanced antibiotic re-






FIG 7 The *dhhP* mutant is defective in cell division. (A) Morphology of 5A4NP1 *dhhP*/pJJ01 grown with IPTG or 5A4NP1 *dhhP*/pJJ01 grown without IPTG examined under a dark-field microscope. Images were taken at a magnification of  $\times 500$ . (B) Quantitation of cell length. A total of 100 randomly selected cells from each strain were measured.

sistance, the conditional *dhhP* mutant of *B. burgdorferi* did not exhibit a dramatic increase in resistance (and showed slightly reduced resistance [ $2\times$ ] to some  $\beta$ -lactam antibiotics) (Table 4). These results suggest that c-di-AMP may function differently in spirochetes than in Gram-positive bacteria.

**The *dhhP* mutant is defective in producing the major virulence factor OspC.** To examine the impact of deletion of *dhhP* on the overall protein profile of *B. burgdorferi*, whole-cell lysates from various strains were harvested and subjected to SDS-PAGE analysis (Fig. 8A). In comparison to the *dhhP* mutant grown in the presence of IPTG, the same mutant in the absence of IPTG had much lower OspC levels. Since *ospC* expression is controlled by RpoS, we further examined the levels of RpoS by immunoblotting. As expected, RpoS was induced in the conditional *dhhP* mutant with IPTG, with an increase of cell density (Fig. 8A). However, RpoS was not detectable in the *dhhP* mutant in the absence of IPTG.

It is well established that expression of *rpoS* is governed by two transcriptional factors, Rrp2 and BosR (63–65). Regulation of Rrp2 occurs at the level of phosphorylation. Since a method to assess the level of Rrp2 phosphorylation within the cell is lacking,

a potential impact of DhhP on Rrp2 phosphorylation could not be examined. Therefore, we focused on regulation of BosR (43, 57, 66). Similar to what was seen with the levels of RpoS, the *dhhP* mutant failed to induce BosR production in the absence of IPTG, indicating that the loss of OspC and RpoS in the *dhhP* mutant was due to a defect in BosR production (Fig. 8A). As expected, the *dhhP* mutant had no *dhhP* mRNA and had a significantly reduced *rpoS* mRNA level (Fig. 8B and C). However, the *bosR* mRNA level was only moderately reduced, with no statistical significance, in the *dhhP* mutant (Fig. 8D), suggesting that deletion of *dhhP* affects BosR mainly at the posttranscriptional level, a phenomenon that has been observed under several other conditions (39, 57, 66). Thus, DhhP appears to be required for BosR accumulation, which in turn controls production of RpoS and RpoS-dependent virulence factors, including OspC.


**DISCUSSION**

Virtually all reported studies on the c-di-AMP signaling thus far have focused on Gram-positive bacteria, with one exception showing that the Gram-negative pathogen *C. trachomatis* is also capable of producing c-di-AMP (27). In this study, we showed that c-di-AMP signaling is also present in the phylum *Spirochaetae*. We demonstrated, *in vitro* and *in vivo*, that the DHH-DHHA1 protein in *B. burgdorferi*, DhhP, functions as a c-di-AMP PDE. Unlike what has been observed in Gram-positive bacteria, DhhP is required for cell growth of *B. burgdorferi*.

The requirement of DhhP for growth of *B. burgdorferi* was surprising, given that none of the c-di-AMP PDEs has been shown to be essential for cell growth, but rather, it has been demonstrated that diadenylate cyclase is indispensable for cell survival in all Gram-positive bacteria examined. Furthermore, the reported PDE-defective mutants (containing increased intracellular c-di-AMP levels) have elevated resistance to peptidoglycan-targeted antibiotics in *S. aureus* (21), *B. subtilis* (15), *S. pyogenes* (26), and *L. monocytogenes* (8). However, deletion of *dhhP* of *B. burgdorferi* did not result in an increased resistance to cell wall-targeting antibiotics (Table 4). Since increased antibiotic resistance in c-di-AMP PDE mutants has been linked to an increased amount of cross-linked peptidoglycan in *S. aureus* (21), we also examined the *dhhP* mutant of *B. burgdorferi* using cryo-electron microscopy in collaboration with Jun Liu at the University of Texas at Houston. We did not observe any significant change in the thickness of the peptidoglycan layer (data not shown). Another phenotype of the *dhhP* mutant that is different from the phenotypes of other PDE mutants in Gram-positive bacteria concerned differences in cell shape. The *dhhP* mutant of *B. burgdorferi* had an increased cell length, whereas inactivation of *gdpP* resulted in a reduced cell size of *S. aureus* (21). In *B. subtilis*, curled nonseparated cell filaments were observed upon cellular accumulation of c-di-AMP (67). All these findings suggest that c-di-AMP may function differently in modulating cell wall homeostasis of *B. burgdorferi*. As spirochetes

TABLE 4 MIC of *B. burgdorferi* strains for various antibiotics

Strain	MIC ( $\mu\text{g/ml}$ ) of:					
	Ampicillin	Aztreonam	Cefuroxime	Cefixime	Vancomycin	Oxacillin
5A4NP1	2.5	25	0.156	0.625	0.625	1.25
5A4NP1 <i>dhhP</i> /pJJ01 +IPTG	2.5	25	0.156	0.625	0.625	1.25
5A4NP1 <i>dhhP</i> /pJJ01 –IPTG	1.25	25	0.156	0.625	0.312	0.625



**FIG 8** The *dhhp* mutant is defective in the production of the major virulence factor OspC. (A) The conditional *dhhp* mutant was inoculated at a higher cell density ( $3 \times 10^5$  cells/ml) in BSK-II medium with or without IPTG at 37°C. Cells were harvested at different days (labeled at the top) and subjected to SDS-PAGE and immunoblotting analyses. Flab served as a loading control. The bands corresponding to OspC are indicated by an arrow. (B to D) qRT-PCR assessment of expression of *dhhp* (B), *rpoS* (C), and *bosR* (D). RNA was extracted from cells harvested at day 3 and subjected to qRT-PCR analyses. Levels of expression of each gene were normalized with *recA*.

have a cell wall structure similar to that of Gram-negative bacteria (with dual membranes), it will be interesting to investigate the roles of c-di-AMP phosphodiesterases in proteobacteria that appear to possess c-di-AMP signaling pathways (5).

Most of the c-di-AMP PDEs reported thus far have been grouped into the GdpP family, including *B. subtilis* YybT (17), *S. aureus* GdpP, *S. pyogenes* GdpP (26), and *S. pneumoniae* Pde1 (7). Besides DHH-DHHA1 domains responsible for c-di-AMP hydrolysis, proteins in the GdpP family contain an additional degenerate GGDEF domain and PAS domain (5). An intact GGDEF domain is responsible for diguanylate cyclase activity involved in synthesis of c-di-GMP (34, 68, 69). The GGDEF domain within GdpP, however, has no such activity but is capable of hydrolyzing ATP (17) and may regulate the activity of the DHH-DHHA1 domains (21). The PAS domain in GdpP has been shown to bind to heme and plays a regulatory role in c-di-AMP PDE's activity (70). These proteins also have transmembrane domains and are associated with the cell membrane (5). In this regard, *B. burgdorferi* DhhP has a single DHH-DHHA1 domain module and lacks transmembrane and regulatory domains. This domain structure suggests that DhhP is a cytoplasmic protein and its activity may be regulated at the level of gene expression, not at the level of its enzymatic activity. In this study, we showed that *dhhp* expression is growth phase dependent, being highest during the exponential phase (Fig. 6).

Given that the *dhhp* mutant has an impaired growth phenotype under the *in vitro* cultivation conditions, it was not surprising that the same mutant was not able to replicate in the dialysis membrane implanted in the rat peritoneal cavity (the DMC conditions) or when directly inoculated into mice. These results suggest that the requirement for DhhP does not depend on the environmental niches where *B. burgdorferi* resides, and elevated levels of c-di-AMP likely inhibit one or more cellular processes that are essential for *B. burgdorferi* growth. One of the cellular processes affected by c-di-AMP may be cell division, as the conditional c-di-AMP mutant showed increased cell length (Fig. 7). The *dhhp* mutant was also unable to produce RpoS and RpoS-dependent virulence factors, such as OspC. This defect could be simply due to its growth defect, as induction of RpoS and OspC is growth rate dependent (71).

This study focused on c-di-AMP PDE of *B. burgdorferi*. Inactivation of *dhhp* resulted in accumulation of cellular c-di-AMP, indicating that *B. burgdorferi* is capable of producing the second messenger c-di-AMP. The enzyme diadenylate cyclase (DAC) for c-di-AMP synthesis contains a conserved DisA\_N domain (21). In this regard, we have found that the hypothetical protein BB0008 in the *B. burgdorferi* genome encodes a putative DisA\_N domain (E value =  $5.45 \times 10^{-34}$ ), which makes BB0008 a strong candidate as a DAC of *B. burgdorferi*. Currently, no information is available about BB0008; its expression is not regulated by temperature, DMC conditions, or the two-component system Hk1-Rrp1 or Hk2-Rrp2 based on the published microarray data (37, 38, 72–79). We are currently in the process of investigating the role of BB0008 in the c-di-AMP signaling of *B. burgdorferi*.

## ACKNOWLEDGMENTS

Funding for this work was partially provided by NIH grants AI083640 and AI085242 and the Indiana INGEN and METACyt grants of Indiana University. This study was funded by the Lilly Endowment, Inc. (to X.F.Y.), by the National Science Foundation of China (81171611 to Y.L. and 31070050 to J.-J.Z.), and by the U.S. National Science Foundation (MCB1052575 to M.G.). This investigation was partially conducted in a facility with support from research facilities improvement program grant number C06 RR015481-01 from the National Center for Research Resources, NIH.

## REFERENCES

- Romling U. 2008. Great times for small molecules: c-di-AMP, a second messenger candidate in Bacteria and Archaea. *Sci. Signal.* 1:pe39. <http://dx.doi.org/10.1126/scisignal.133pe39>.
- Gomelsky M. 2011. cAMP, c-di-GMP, c-di-AMP and now cGMP: bacteria use them all! *Mol. Microbiol.* 79:562–565. <http://dx.doi.org/10.1111/j.1365-2958.2010.07514.x>.
- McDonough KA, Rodriguez A. 2012. The myriad roles of cyclic AMP in microbial pathogens: from signal to sword. *Nat. Rev. Microbiol.* 10:27–38. <http://dx.doi.org/10.1038/nrmicro2688>.
- Kalia D, Meray G, Nakayama S, Zheng Y, Zhou J, Luo Y, Guo M, Roembke BT, Sintim HO. 2013. Nucleotide, c-di-GMP, c-di-AMP, cGMP, cAMP, (p)ppGpp signaling in bacteria and implications in pathogenesis. *Chem. Soc. Rev.* 42:305–341. <http://dx.doi.org/10.1039/c2cs35206k>.
- Corrigan RM, Grundling A. 2013. Cyclic di-AMP: another second messenger enters the fray. *Nat. Rev. Microbiol.* 11:513–524. <http://dx.doi.org/10.1038/nrmicro3069>.



6. Witte G, Hartung S, Buttner K, Hopfner KP. 2008. Structural biochemistry of a bacterial checkpoint protein reveals diadenylate cyclase activity regulated by DNA recombination intermediates. *Mol. Cell* 30:167–178. <http://dx.doi.org/10.1016/j.molcel.2008.02.020>.
7. Bai Y, Yang J, Eisele LE, Underwood AJ, Koestler BJ, Waters CM, Metzger DW, Bai G. 2013. Two DHH subfamily 1 proteins in *Streptococcus pneumoniae* possess cyclic di-AMP phosphodiesterase activity and affect bacterial growth and virulence. *J. Bacteriol.* 195:5123–5132. <http://dx.doi.org/10.1128/JB.00769-13>.
8. Witte CE, Whiteley AT, Burke TP, Sauer JD, Portnoy DA, Woodward JJ. 2013. Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall homeostasis, and establishment of infection. *mBio* 4:e00282–13. <http://dx.doi.org/10.1128/mBio.00282-13>.
9. French CT, Lao P, Loraine AE, Matthews BT, Yu H, Dybvig K. 2008. Large-scale transposon mutagenesis of *Mycoplasma pulmonis*. *Mol. Microbiol.* 69:67–76. <http://dx.doi.org/10.1111/j.1365-2958.2008.06262.x>.
10. Woodward JJ, Iavarone AT, Portnoy DA. 2010. c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science* 328:1703–1705. <http://dx.doi.org/10.1126/science.1189801>.
11. Kamegaya T, Kuroda K, Hayakawa Y. 2011. Identification of a *Streptococcus pyogenes* SF370 gene involved in production of c-di-AMP. *Nagoya J. Med. Sci.* 73:49–57. [http://ir.nul.nagoya-u.ac.jp/jspui/bitstream/2237/14915/1/v73n12p49\\_57.pdf](http://ir.nul.nagoya-u.ac.jp/jspui/bitstream/2237/14915/1/v73n12p49_57.pdf).
12. Song JH, Ko KS, Lee JY, Baek JY, Oh WS, Yoon HS, Jeong JY, Chun J. 2005. Identification of essential genes in *Streptococcus pneumoniae* by allelic replacement mutagenesis. *Mol. Cells* 19:365–374. [http://chunlab.snu.ac.kr:8080/Plone/publications/Song\\_2005\\_MAC.pdf](http://chunlab.snu.ac.kr:8080/Plone/publications/Song_2005_MAC.pdf).
13. Glass JI, Assad-Garcia N, Alperovich N, Yooseph S, Lewis MR, Maruf M, Hutchison CA, III, Smith HO, Venter JC. 2006. Essential genes of a minimal bacterium. *Proc. Natl. Acad. Sci. U. S. A.* 103:425–430. <http://dx.doi.org/10.1073/pnas.0510013103>.
14. Chaudhuri RR, Allen AG, Owen PJ, Shalom G, Stone K, Harrison M, Burgis TA, Lockyer M, Garcia-Lara J, Foster SJ, Pleasance SJ, Peters SE, Maskell DJ, Charles IG. 2009. Comprehensive identification of essential *Staphylococcus aureus* genes using transposon-mediated differential hybridisation (TMDH). *BMC Genomics* 10:291. <http://dx.doi.org/10.1186/1471-2164-10-291>.
15. Luo Y, Helmann JD. 2012. Analysis of the role of *Bacillus subtilis* s<sup>M</sup> in  $\beta$ -lactam resistance reveals an essential role for c-di-AMP in peptidoglycan homeostasis. *Mol. Microbiol.* 83:623–639. <http://dx.doi.org/10.1111/j.1365-2958.2011.07953.x>.
16. Smith WM, Pham TH, Lei L, Dou JC, Soomro AH, Beatson SA, Dykes GA, Turner MS. 2012. Heat resistance and salt hypersensitivity in *Lactococcus lactis* due to spontaneous mutation of *llmg\_1816* (*gdpP*) induced by high-temperature growth. *Appl. Environ. Microbiol.* 78:7753–7759. <http://dx.doi.org/10.1128/AEM.02316-12>.
17. Rao F, See RY, Zhang DW, Toh DC, Ji Q, Liang ZX. 2010. YybT is a signaling protein that contains a cyclic dinucleotide phosphodiesterase domain and a GGDEF domain with ATPase activity. *J. Biol. Chem.* 285:473–482. <http://dx.doi.org/10.1074/jbc.M109.040238>.
18. Griffiths JM, O'Neill AJ. 2012. Loss of function of the *gdpP* protein leads to joint  $\beta$ -lactam/glycopeptide tolerance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 56:579–581. <http://dx.doi.org/10.1128/AAC.05148-11>.
19. Henry JT, Crosson S. 2011. Ligand-binding PAS domains in a genomic, cellular, and structural context. *Annu. Rev. Microbiol.* 65:261–286. <http://dx.doi.org/10.1146/annurev-micro-121809-151631>.
20. Ponting CP, Aravind L. 1997. PAS: a multifunctional domain family comes to light. *Curr. Biol.* 7:R674–R677. [http://dx.doi.org/10.1016/S0960-9822\(06\)00352-6](http://dx.doi.org/10.1016/S0960-9822(06)00352-6).
21. Corrigan RM, Abbott JC, Burhenne H, Kaever V, Grundling A. 2011. c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. *PLoS Pathog.* 7:e1002217.
22. Cron LE, Stol K, Burghout P, van Selm S, Simonetti ER, Bootsma HJ, Hermans PWM. 2012. Two DHH subfamily 1 proteins contribute to pneumococcal virulence and confer protection against pneumococcal disease. *Infect. Immun.* 80:2974. <http://dx.doi.org/10.1128/IAI.00519-12>.
23. Zhang L, Li W, He ZG. 2013. DarR, a TetR-like transcriptional factor, is a cyclic di-AMP-responsive repressor in *Mycobacterium smegmatis*. *J. Biol. Chem.* 288:3085–3096. <http://dx.doi.org/10.1074/jbc.M112.428110>.
24. Bai Y, Yang J, Zarrella TM, Zhang Y, Metzger DW, Bai G. 22 November 2013. Cyclic di-AMP impairs potassium uptake mediated by a c-di-AMP binding protein in *Streptococcus pneumoniae*. *J. Bacteriol.* <http://dx.doi.org/10.1128/JB.01041-13>.
25. Corrigan RM, Campeotto I, Jegannathan T, Roelofs KG, Lee VT, Gründling A. 2013. Systematic identification of conserved bacterial c-di-AMP receptor proteins. *Proc. Natl. Acad. Sci. U. S. A.* 110:9084–9089. <http://dx.doi.org/10.1073/pnas.1300595110>.
26. Cho KH, Kang SO. 2013. *Streptococcus pyogenes* c-di-AMP phosphodiesterase, GdpP, influences SpeB processing and virulence. *PLoS One* 8:e69425. <http://dx.doi.org/10.1371/journal.pone.0069425>.
27. Barker JR, Koestler BJ, Carpenter VK, Burdette DL, Waters CM, Vance RE, Valdivia RH. 2013. STING-dependent recognition of cyclic di-AMP mediates type I interferon responses during *Chlamydia trachomatis* infection. *mBio* 4:e00018–13. <http://dx.doi.org/10.1128/mBio.00018-13>.
28. Steere AC, Coburn J, Glickstein L. 2004. The emergence of Lyme disease. *J. Clin. Invest.* 113:1093–1101. <http://dx.doi.org/10.1172/JCI21681>.
29. Samuels DS. 2011. Gene regulation in *Borrelia burgdorferi*. *Annu. Rev. Microbiol.* 65:479–499. <http://dx.doi.org/10.1146/annurev.micro.112408.134040>.
30. Radolf JD, Caimano MJ, Stevenson B, Hu LT. 2012. Of ticks, mice and men: understanding the dual-host lifestyle of Lyme disease spirochaetes. *Nat. Rev. Microbiol.* 10:87–99. <http://dx.doi.org/10.1038/nrmicro2714>.
31. Rosa PA, Tilly K, Stewart PE. 2005. The burgeoning molecular genetics of the Lyme disease spirochaete. *Nat. Rev. Microbiol.* 3:129–143. <http://dx.doi.org/10.1038/nrmicro1086>.
32. Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD, Richardson D, Peterson J, Kervlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Wattney L, McDonald L, Artiach P, Bowman C, Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. 1997. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature* 390:580–586. <http://dx.doi.org/10.1038/37551>.
33. Römmling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial second messenger. *Microbiol. Mol. Biol. Rev.* 77:1–52. <http://dx.doi.org/10.1128/MMBR.00043-12>.
34. Hengge R. 2009. Principles of c-di-GMP signalling in bacteria. *Nat. Rev. Microbiol.* 7:263–273. <http://dx.doi.org/10.1038/nrmicro2109>.
35. Jenal U, Malone J. 2006. Mechanisms of cyclic-di-GMP signaling in bacteria. *Annu. Rev. Genet.* 40:385–407. <http://dx.doi.org/10.1146/annurev.genet.40.110405.090423>.
36. Wolfe AJ, Visick KL. 2010. The second messenger cyclic di-GMP. ASM Press, Washington, DC.
37. Caimano MJ, Kenedy MR, Kairu T, Desrosiers DC, Harman M, Dunham-Ems S, Akins DR, Pal U, Radolf JD. 2011. The hybrid histidine kinase Hk1 is part of a two-component system that is essential for survival of *Borrelia burgdorferi* in feeding *Ixodes scapularis* ticks. *Infect. Immun.* 79:3117–3130. <http://dx.doi.org/10.1128/IAI.05136-11>.
38. He M, Ouyang Z, Troxell B, Xu H, Moh A, Piesman J, Norgard MV, Gomelsky M, Yang XF. 2011. Cyclic di-GMP is essential for the survival of the Lyme disease spirochete in ticks. *PLoS Pathog.* 7:e1002133. <http://dx.doi.org/10.1371/journal.ppat.1002133>.
39. He M, Zhang J-J, Ye M, Lou Y, Yang XF. 11 November 2013. The cyclic di-GMP receptor PlzA controls virulence gene expression through RpoS in *Borrelia burgdorferi*. *Infect. Immun.* <http://dx.doi.org/10.1128/IAI.01238-13>.
40. Kostick JL, Szkotnicki LT, Rogers EA, Bocci P, Raffaelli N, Marconi RT. 2011. The diguanylate cyclase, Rrp1, regulates critical steps in the enzootic cycle of the Lyme disease spirochetes. *Mol. Microbiol.* 81:219–231. <http://dx.doi.org/10.1111/j.1365-2958.2011.07687.x>.
41. Sultan SZ, Pitzer JE, Boquai T, Hobbs G, Miller MR, Motaleb MA. 2011. Analysis of the HD-GYP domain cyclic dimeric GMP phosphodiesterase reveals a role in motility and the enzootic life cycle of *Borrelia burgdorferi*. *Infect. Immun.* 79:3273–3283. <http://dx.doi.org/10.1128/IAI.05153-11>.
42. Sultan SZ, Pitzer JE, Miller MR, Motaleb MA. 2010. Analysis of a *Borrelia burgdorferi* phosphodiesterase demonstrates a role for cyclic-diguanylate monophosphate in motility and virulence. *Mol. Microbiol.* 77:128–142. <http://dx.doi.org/10.1111/j.1365-2958.2010.07191.x>.
43. Sze CW, Smith A, Choi YH, Yang X, Pal U, Yu A, Li C. 2013. Study of the response regulator Rrp1 reveals its regulatory role in chitinase utilization and virulence of *Borrelia burgdorferi*. *Infect. Immun.* 81:1775–1787. <http://dx.doi.org/10.1128/IAI.00050-13>.
44. Kawabata H, Norris SJ, Watanabe H. 2004. BBE02 disruption mutants of *Borrelia burgdorferi* B31 have a highly transformable, infectious pheno-

- type. *Infect. Immun.* 72:7147–7154. <http://dx.doi.org/10.1128/IAI.72.12.7147-7154.2004>.
45. Barbour AG. 1984. Isolation and cultivation of Lyme disease spirochetes. *Yale J. Biol. Med.* 57:521–525.
  46. Fang X, Gomelsky M. 2010. A post-translational, c-di-GMP-dependent mechanism regulating flagellar motility. *Mol. Microbiol.* 76:1295–1305. <http://dx.doi.org/10.1111/j.1365-2958.2010.07179.x>.
  47. Ryjenkov DA, Simm R, Romling U, Gomelsky M. 2006. The PilZ domain is a receptor for the second messenger c-di-GMP: the PilZ domain protein YcgR controls motility in enterobacteria. *J. Biol. Chem.* 281:30310–30314. <http://dx.doi.org/10.1074/jbc.C600179200>.
  48. Bobrov AG, Kirillina O, Ryjenkov DA, Waters CM, Price PA, Fetherston JD, Mack D, Goldman WE, Gomelsky M, Perry RD. 2011. Systematic analysis of cyclic di-GMP signalling enzymes and their role in biofilm formation and virulence in *Yersinia pestis*. *Mol. Microbiol.* 79:533–551. <http://dx.doi.org/10.1111/j.1365-2958.2010.07470.x>.
  49. Charon NW, Cockburn A, Li C, Liu J, Miller KA, Miller MR, Motaleb MA, Wolgemuth CW. 2012. The unique paradigm of spirochete motility and chemotaxis. *Annu. Rev. Microbiol.* 66:349–370. <http://dx.doi.org/10.1146/annurev-micro-092611-150145>.
  50. Blevins JS, Revel AT, Smith AH, Bachlani GN, Norgard MV. 2007. Adaptation of a luciferase gene reporter and *lac* expression system to *Borrelia burgdorferi*. *Appl. Environ. Microbiol.* 73:1501–1513. <http://dx.doi.org/10.1128/AEM.02454-06>.
  51. Chen Y, Holcomb C, Moore H. 1993. Expression and localization of two low molecular weight GTP-binding proteins, Rab8 and Rab10, by epitope tag. *Proc. Natl. Acad. Sci. U. S. A.* 90:6508–6512. <http://dx.doi.org/10.1073/pnas.90.14.6508>.
  52. Yang XF, Pal U, Alani SM, Fikrig E, Norgard MV. 2004. Essential role for OspA/B in the life cycle of the Lyme disease spirochete. *J. Exp. Med.* 199:641–648. <http://dx.doi.org/10.1084/jem.20031960>.
  53. Samuels DS. 1995. Electrotransformation of the spirochete *Borrelia burgdorferi*. *Methods Mol. Biol.* 47:253–259.
  54. Elias AF, Bono JL, Kupko JJ, Stewart PE, Krum JG, Rosa PA. 2003. New antibiotic resistance cassettes suitable for genetic studies in *Borrelia burgdorferi*. *J. Mol. Microbiol. Biotechnol.* 6:29–40. <http://dx.doi.org/10.1159/000073406>.
  55. Bunikis I, Kutschan-Bunikis S, Bonde M, Bergström S. 2011. Multiplex PCR as a tool for validating plasmid content of *Borrelia burgdorferi*. *J. Microbiol. Methods* 86:243–247. <http://dx.doi.org/10.1016/j.mimet.2011.05.004>.
  56. Dever LL, Jorgensen JH, Barbour AG. 1992. In vitro antimicrobial susceptibility testing of *Borrelia burgdorferi*: a microdilution MIC method and time-kill studies. *J. Clin. Microbiol.* 30:2692–2697.
  57. Troxell B, Ye M, Yang Y, Carrasco SE, Lou Y, Yang XF. 2013. Manganese and zinc regulate virulence determinants in *Borrelia burgdorferi*. *Infect. Immun.* 81:2743–2752. <http://dx.doi.org/10.1128/IAI.00507-13>.
  58. Xu HJ, He M, He JJ, Yang XF. 2010. Role of the surface lipoprotein BBA07 in the enzootic cycle of *Borrelia burgdorferi*. *Infect. Immun.* 78:2910–2918. <http://dx.doi.org/10.1128/IAI.00372-10>.
  59. He M, Oman T, Xu H, Blevins JS, Norgard MV, Yang XF. 2008. Abrogation of *ospAB* constitutively activates the Rrp2-RpoN-RpoS pathway (sigmaN-sigmaS cascade) in *Borrelia burgdorferi*. *Mol. Microbiol.* 70:1453–1464. <http://dx.doi.org/10.1111/j.1365-2958.2008.06491.x>.
  60. Akins DR, Bourell KW, Caimano MJ, Norgard MV, Radolf JD. 1998. A new animal model for studying Lyme disease spirochetes in a mammalian host-adapted state. *J. Clin. Invest.* 101:2240–2250. <http://dx.doi.org/10.1172/JCI2325>.
  61. Caimano M. 2005. Cultivation of *Borrelia burgdorferi* in dialysis membrane chambers in rat peritonea. *Curr. Protoc. Microbiol.* 2005:12.C.3.1-12.C.3.13. <http://dx.doi.org/10.1002/9780471729259.mc12c03s00>.
  62. Gilbert MA, Morton EA, Bundle SF, Samuels DS. 2007. Artificial regulation of *ospC* expression in *Borrelia burgdorferi*. *Mol. Microbiol.* 63:1259–1273. <http://dx.doi.org/10.1111/j.1365-2958.2007.05593.x>.
  63. Ouyang Z, Kumar M, Kariu T, Haq S, Goldberg M, Pal U, Norgard MV. 2009. BosR (BB0647) governs virulence expression in *Borrelia burgdorferi*. *Mol. Microbiol.* 74:1331–1343. <http://dx.doi.org/10.1111/j.1365-2958.2009.06945.x>.
  64. Hyde JA, Shaw DK, Smith Iii R, Trzeciakowski JP, Skare JT. 2009. The BosR regulatory protein of *Borrelia burgdorferi* interfaces with the RpoS regulatory pathway and modulates both the oxidative stress response and pathogenic properties of the Lyme disease spirochete. *Mol. Microbiol.* 74:1344–1355. <http://dx.doi.org/10.1111/j.1365-2958.2009.06951.x>.
  65. Yang XF, Alani SM, Norgard MV. 2003. The response regulator Rrp2 is essential for the expression of major membrane lipoproteins in *Borrelia burgdorferi*. *Proc. Natl. Acad. Sci. U. S. A.* 100:11001–11006. <http://dx.doi.org/10.1073/pnas.1834315100>.
  66. Hyde JA, Trzeciakowski JP, Skare JT. 2007. *Borrelia burgdorferi* alters its gene expression and antigenic profile in response to CO<sub>2</sub> levels. *J. Bacteriol.* 189:437–445. <http://dx.doi.org/10.1128/JB.01109-06>.
  67. Mehne FM, Gunka K, Eilers H, Herzberg C, Kaever V, Stulke J. 2013. Cyclic di-AMP homeostasis in *Bacillus subtilis*: both lack and high-level accumulation of the nucleotide are detrimental for cell growth. *J. Biol. Chem.* 288:2004–2017. <http://dx.doi.org/10.1074/jbc.M112.395491>.
  68. Ausmees N, Mayer R, Weinhouse H, Volman G, Amikam D, Benziman M, Lindberg M. 2001. Genetic data indicate that proteins containing the GGDEF domain possess diguanylate cyclase activity. *FEMS Microbiol. Lett.* 204:163–167. <http://dx.doi.org/10.1111/j.1574-6968.2001.tb10880.x>.
  69. Ryjenkov DA, Tarutina M, Moskvina OV, Gomelsky M. 2005. Cyclic diguanylate is a ubiquitous signaling molecule in bacteria: insights into biochemistry of the GGDEF protein domain. *J. Bacteriol.* 187:1792–1798. <http://dx.doi.org/10.1128/JB.187.5.1792-1798.2005>.
  70. Rao F, Ji Q, Soehano I, Liang ZX. 2011. Unusual heme-binding PAS domain from YybT family proteins. *J. Bacteriol.* 193:1543–1551. <http://dx.doi.org/10.1128/JB.01364-10>.
  71. Jutras BL, Chenail AM, Stevenson B. 2013. Changes in bacterial growth rate govern expression of the *Borrelia burgdorferi* OspC and Erp infection-associated surface proteins. *J. Bacteriol.* 195:757–764. <http://dx.doi.org/10.1128/JB.01956-12>.
  72. Freedman JC, Rogers EA, Kostick JL, Zhang H, Iyer R, Schwartz I, Marconi RT. 2010. Identification and molecular characterization of a cyclic-di-GMP effector protein, PlzA (BB0733): additional evidence for the existence of a functional cyclic-di-GMP regulatory network in the Lyme disease spirochete, *Borrelia burgdorferi*. *FEMS Immunol. Med. Microbiol.* 58:285–294. <http://dx.doi.org/10.1111/j.1574-695X.2009.00635.x>.
  73. Boardman BK, He M, Ouyang Z, Xu H, Pang X, Yang XF. 2008. Essential role of the response regulator Rrp2 in the infectious cycle of *Borrelia burgdorferi*. *Infect. Immun.* 76:3844–3853. <http://dx.doi.org/10.1128/IAI.00467-08>.
  74. Brooks CS, Hefty PS, Jolliffe SE, Akins DR. 2003. Global analysis of *Borrelia burgdorferi* genes regulated by mammalian host-specific signals. *Infect. Immun.* 71:3371–3383. <http://dx.doi.org/10.1128/IAI.71.6.3371-3383.2003>.
  75. Caimano M, Iyer R, Eggers C, Gonzalez C, Morton E, Gilbert M, Schwartz I, Radolf J. 2007. Analysis of the RpoS regulon in *Borrelia burgdorferi* in response to mammalian host signals provides insight into RpoS function during the enzootic cycle. *Mol. Microbiol.* 65:1193–1217. <http://dx.doi.org/10.1111/j.1365-2958.2007.05860.x>.
  76. Fisher MA, Grimm D, Henion AK, Elias AF, Stewart PE, Rosa PA, Gherardini FC. 2005. *Borrelia burgdorferi*  $\sigma^{54}$  is required for mammalian infection and vector transmission but not for tick colonization. *Proc. Natl. Acad. Sci. U. S. A.* 102:5162–5167. <http://dx.doi.org/10.1073/pnas.0408536102>.
  77. Ojaimi C, Brooks C, Casjens S, Rosa P, Elias A, Barbour A, Jasinskas A, Benach J, Katona L, Radolf J, Caimano M, Skare J, Swingle K, Akins D, Schwartz I. 2003. Profiling of temperature-induced changes in *Borrelia burgdorferi* gene expression by using whole genome arrays. *Infect. Immun.* 71:1689–1705. <http://dx.doi.org/10.1128/IAI.71.4.1689-1705.2003>.
  78. Ouyang Z, Blevins JS, Norgard MV. 2008. Transcriptional interplay among the regulators Rrp2, RpoN, and RpoS in *Borrelia burgdorferi*. *Microbiology* 154:2641–2658. <http://dx.doi.org/10.1099/mic.0.2008/019992-0>.
  79. Revel AT, Talaat AM, Norgard MV. 2002. DNA microarray analysis of differential gene expression in *Borrelia burgdorferi*, the Lyme disease spirochete. *Proc. Natl. Acad. Sci. U. S. A.* 99:1562–1567. <http://dx.doi.org/10.1073/pnas.032667699>.